arXiv:1208.4510v1 [math.DS] 22 Aug 2012

FIXED POINTS OF NILPOTENT ACTIONS ON $\mathbb{S}^2$

JAVIER RIBÓN

Abstract

We prove that a nilpotent subgroup of orientation preserving C^1 diffeomorphisms of $\mathbb{S}^2$ has a finite orbit of period at most two. We also prove that a nilpotent subgroup of orientation preserving C^1 diffeomorphisms of $\mathbb{R}^2$ preserving a compact set has a global fixed point. These results generalize theorems of Franks, Handel and Parwani for the abelian case.

We show that a nilpotent subgroup of orientation preserving C^1 diffeomorphisms of $\mathbb{S}^2$ that has an odd finite orbit also has a global fixed point. Moreover we study the properties of the period two orbits of nilpotent fixed-point-free subgroups of orientation preserving C^1 diffeomorphisms of $\mathbb{S}^2$.

1. Introduction

Consider a closed surface S with non-vanishing Euler characteristic. The classical Poincaré theorem asserts that any C^1 vector field defined in S has a singular point. Lima [16] proved that this result can be extended to any finite family of pairwise commuting C^1 vector fields and then to any continuous action of a connected abelian finite-dimensional Lie group G on S. Plante generalized the previous result for connected nilpotent finite-dimensional Lie groups [18].

If the action is not necessarily connected and continuous the natural generalization is studying fixed points for subgroups of $\mathrm{Diff}_0^1(S)$ (the group of isotopic to the identity C^1 diffeomorphisms of S). The Lefschetz fixed point formula implies that any element of $\mathrm{Diff}_0^1(S)$ has a fixed point. An intermediate case between the study of continuous actions of Lie algebras and the general case is obtained by considering subgroups G of $\mathrm{Diff}^1(S)$ whose generators are close to the identity map in the C^1 topology. The orbits of $\phi \in G$ are contained in poligonal

---

e-mail address: javier@mat.uff.br.

MSC-class. Primary: 37C85, 37E30; Secondary: 37C25, 55M20, 57S25.

Keywords: local diffeomorphism, derived length, solvable group, nilpotent group.

curves that are almost tangent to the trajectories of a vector field X associated to ϕ (see [4]). Bonatti took profit of this flow-like behavior to prove that any commutative subgroup G of $\mathrm{Diff}_0^1(S)$ $(\chi(S) \neq 0)$ whose generators are C^1-close to the identity has a global fixed point [3] [4]. This result was generalized by Druck, Fang and Firmo for nilpotent groups of diffeomorphisms of the sphere in the C^1-close to the identity setting [7].

A natural idea is trying to extend the previous results to general subgroups of $\mathrm{Diff}_0^1(\mathbb{S}^2)$ by replacing flows (or almost flows) with isotopies. In this way Franks, Handel and Parwani solve completely the problem of existence of fixed points for commutative groups. They prove that an abelian subgroup of $\mathrm{Diff}_0^1(S)$ $(\chi(S) < 0)$ has always global fixed points (and even more that there are fixed points in the identity lift of the group) [8]. In the case $\chi(S) > 0$, i.e. $S = \mathbb{S}^2$, the group $\mathrm{Diff}_0^1(\mathbb{S}^2)$ coincides with the group $\mathrm{Diff}_+^1(\mathbb{S}^2)$ of orientation-preserving diffeomorphisms. In contrast with the other cases there exist fixed-point-free commutative groups of Möbius transformations (the group generated by two involutions with disjoint fixed point sets such that any involution permutes the fixed points of the other involution). Franks, Handel and Parwani characterize the subgroups of $\mathrm{Diff}_+^1(\mathbb{S}^2)$ that have global fixed points.

Theorem 1.1. [9] *For any abelian subgroup G of $\mathrm{Diff}_+^1(\mathbb{S}^2)$ there is a subgroup H of index at most two such that $\mathrm{Fix}(H) \neq \emptyset$. Moreover, $\mathrm{Fix}(G) \neq \emptyset$ if and only if $w(\phi, \eta) = 0$ for all $\phi, \eta \in G$.*

The invariant $w(\phi, \eta)$ is defined for commuting elements of $\mathrm{Homeo}_+(\mathbb{S}^2)$ and belongs to $\mathbb{Z}/2\mathbb{Z}$ (see subsection 5.1).

We generalize Theorem 1.1 for nilpotent groups.

Theorem 1.2. *Let $G \subset \mathrm{Diff}_+^1(\mathbb{S}^2)$ be a nilpotent group. Then G has an orbit of period at most 2.*

Let us remark that the existence of periodic orbits for nilpotent groups of real analytic diffeomorphisms of the sphere was already known (Ghys [11]).

The proof of Theorem 1.1 relies on the study of lifts of the group to universal coverings of the complementary of some closed invariant sets and a fixed point lemma whose generalization for the nilpotent case follows.

Theorem 1.3. *Let $G \subset \mathrm{Diff}_+^1(\mathbb{R}^2)$ be a finitely generated nilpotent group that preserves a non-empty compact set. Then $\mathrm{Fix}(G)$ is not empty.*

The following results are applications of Theorem 1.2.

Corollary 1.1. *If G is a nilpotent subgroup of $\mathrm{Diff}_+^1(\mathbb{D}^2)$ then $\mathrm{Fix}(G)$ is non-empty.*

Theorem 1.4. *(Generalization of Cartwright-Littlewood theorem). Let $G \subset \mathrm{Diff}_+^1(\mathbb{R}^2)$ be a nilpotent group preserving a full continuum K of the plane. Then G has a global fixed point in K.*

Let G be a nilpotent subgroup of $\mathrm{Diff}_+^1(\mathbb{S}^2)$. In general it is not possible to define an invariant $w : G \times G \to \mathbb{Z}/2\mathbb{Z}$ such that w is symmetric, it is a morphism of groups in every component and satisfies that the subgroups H of G with $w_{|H \times H} \equiv 0$ are exactly the subgroups of G that have global fixed points. The group $< 1/z, iz > \subset \mathrm{PGL}(2, \mathbb{C})$ is an example. It is isomorphic to the dihedral group D_4. We denote $D_2 =< 1/z, -z >$; if the invariant exists then $w_{|D_2 \times D_2} \equiv 0$ since $w(1/z, -z) = 2w(1/z, iz) = 0$ but D_2 is fixed-point-free. Anyway we can still prove the following parity theorem:

Theorem 1.5. *Let $G \subset \mathrm{Diff}_+^1(\mathbb{S}^2)$ be a nilpotent group. Suppose that there exists a finite G-invariant set F such that $\sharp F$ is odd. Then $\mathrm{Fix}(G)$ is not empty.*

The previous theorem can be also interpreted as a result on fixed-point-free groups. Any finite orbit of a fixed-point-free nilpotent subgroup of $\mathrm{Diff}_+^1(\mathbb{S}^2)$ has an even number of elements. We study the properties of these groups in section 5. We say that a finite orbit of G with p elements is a p-orbit of G. Consider 2-orbits $\mathcal{O}_1$ and $\mathcal{O}_2$ of a subgroup G of $\mathrm{Homeo}_+(\mathbb{S}^2)$. We say that $\mathcal{O}_1$ and $\mathcal{O}_2$ belong to the same class if we have

$$\phi_{|\mathcal{O}_1} \equiv Id \Leftrightarrow \phi_{|\mathcal{O}_2} \equiv Id$$

for any $\phi \in G$. Curiously fixed-point-free nilpotent subgroups of $\mathrm{Diff}_+^1(\mathbb{S}^2)$ also have few 2-orbits.

Theorem 1.6. *(Scarcity of 2-orbits). Let G be a fixed-point-free nilpotent subgroup of $\mathrm{Diff}_+^1(\mathbb{S}^2)$. Then the number of classes of 2-orbits is either 1 or 3. Moreover, if G is commutative it is 3.*

2. Technical setup

Let us introduce some of the tools that we are going to use throughout the paper.

2.1. **Groups.** The main results are proved by induction on the class of nilpotency k of G.

Definition 2.1. Let G be a group. We denote by $Z^{(l)}(G)$ the l^{th} member of the upper central series of G. If G is nilpotent then $c = \min\{k \in \mathbb{N} \cup \{0\} : Z^{(k)}(G) = G\}$ is the nilpotency class of G.

Definition 2.2. Let G be a nilpotent group. We say that G is in $\mathcal{G}_{k,p}$ if G is a nilpotent group of nilpotency class $\leq k+1$ such that $G =< Z^{(k)}(G), \alpha_1, \ldots, \alpha_p >$ for some $\alpha_1, \ldots, \alpha_p \in G$.

Remark 2.1. Given a group $G =< Z^{(k)}(G), \alpha_1, \ldots, \alpha_p >$ the group $< Z^{(k)}(G), \alpha_1, \ldots, \alpha_{p-1} >$ belongs to $\mathcal{G}_{k,p-1}$.

Proposition 2.1. *(see page 20* [20]*) Let G be a group acting by isometries on a finite tree. Then the action has a fixed point.*

2.2. **Nielsen classes.** A frequent trick in this paper is to consider a lift $\tilde{G}$ of $G \subset \mathrm{Diff}_+^1(\mathbb{S}^2)$, deducing properties of $\tilde{G}$ and interpreting them in terms of G.

Definition 2.3. Let $G \subset \mathrm{Homeo}_+(S)$ be a group defined in a manifold S. We say that $\tilde{G} \subset \mathrm{Homeo}_+(M)$ is a lift if

- M is the universal covering of a G-invariant open set U of S. We denote by $\pi : M \to U$ the covering transformation.
- There exists a unique $\tilde{\phi} \in \tilde{G}$ such that $\phi \circ \pi = \pi \circ \tilde{\phi}$ for any $\phi \in G$. Reciprocally given $\psi \in \tilde{G}$ there exists $\phi \in G$ such that $\phi \circ \pi = \pi \circ \psi$.

The group $G_{|U}$ induced by G by restriction to U is isomorphic to $\tilde{G}$. It is clear that $\mathrm{Fix}(\tilde{G}) \neq \emptyset$ implies $\mathrm{Fix}(G) \cap U \neq \emptyset$.

We use Nielsen classes to analyze the properties of the fixed point sets of the elements of a lift.

Definition 2.4. Let $\phi \in \mathrm{Homeo}_+(S)$. Consider a compact ϕ-invariant set C. Let $x, y \in \mathrm{Fix}(\phi) \setminus C$. We say that x is *Nielsen equivalent to y relative to C* if there exists an arc ρ from x to y such that ρ and $\phi(\rho)$ are homotopic relative to endpoints in $S \setminus C$.

Definition 2.5. Let $\phi \in \mathrm{Homeo}_+(S)$. Consider a compact ϕ-invariant set C. Let $x \in \mathrm{Fix}(\phi) \setminus C$. Consider the connected component N of $S \setminus C$ containing x and its universal covering $\tilde{N}$. We say that $\tilde{\phi} : \tilde{N} \to \tilde{N}$ is a lift of ϕ for the Nielsen class of x if $\mathrm{Fix}(\tilde{\phi})$ contains a lift of x.

Notice that x and y are Nielsen equivalent if they belong to the same connected component N of $S \setminus C$ and there exists some lift (and hence every lift) $\tilde{\phi} : \tilde{N} \to \tilde{N}$ of ϕ for the Nielsen class of x that also fixes a lift of y.

The next definitions are used in Proposition 2.3 (which is indeed the Proposition 4.2 of [9]). They are required to relate properties of the images of a group $G \subset \mathrm{Diff}^1_+(\mathbb{S}^2)$ in the group of classes of homeomorphisms modulo isotopy (relative to some compact G-invariant set) with properties of G.

Definition 2.6. Let η_t be an isotopy rel C from $\eta_0 = \phi$ to η_1. Consider $x \in \mathrm{Fix}(\phi) \setminus C$ and a lift $\tilde{\phi}$ of ϕ for the Nielsen class of x. There exists a unique continuous lift $\tilde{\eta}_t$ of η_t such that $\tilde{\eta}_0 = \tilde{\phi}$. *The Nielsen class determined by the Nielsen class of* x *and* η_t is by definition the Nielsen class determined by $\tilde{\eta}_t$, i.e. the elements of the Nielsen class are the projections of elements of $\mathrm{Fix}(\tilde{\eta}_t)$.

Definition 2.7. We say that the Nielsen class of $x \in \mathrm{Fix}(\phi) \setminus C$ is compactly covered if there exists some lift $\tilde{\phi} : \tilde{N} \to \tilde{N}$ (and hence every lift) of ϕ for the Nielsen class of x such that $\mathrm{Fix}(\tilde{\phi})$ is compact.

Definition 2.8. Suppose that P is an isolated puncture of N. We say that the Nielsen class of $x \in \mathrm{Fix}(\phi) \setminus C$ *peripherally contains* P if there is a properly immersed ray $\rho \subset N$ based at x that converges to P such that $\phi(\rho)$ is properly homotopic to ρ relative to x.

Definition 2.9. Let $\phi \in \mathrm{Homeo}_+(\mathbb{R}^2)$. Suppose that $C \subset \mathbb{R}^2$ is a non-empty compact ϕ-invariant set with $C \cap \mathrm{Fix}(\phi) = \emptyset$. We define $P(C, \phi)$ as the union of the Nielsen classes of $\mathrm{Fix}(\phi)$ rel C which do not peripherally contain ∞.

Periodic points of orientation-preserving homeomorphisms of $\mathbb{R}^2$ turn around fixed points [10] [15]. The next result is a generalization of such phenomenon for compact invariant sets with no fixed points.

Proposition 2.2. *(Prop. 5.3* [9]*) Let* $\phi \in \mathrm{Homeo}_+(\mathbb{R}^2)$*. Suppose that* $C \subset \mathbb{R}^2$ *is a non-empty compact* ϕ*-invariant set with* $C \cap \mathrm{Fix}(\phi) = \emptyset$*. Then* $P(C, \phi)$ *is a non-empty compact set.*

2.3. **Realization of subgroups of the mapping class group.** An idea in fixed point theory is that irreducible models of elements or groups contain the minimum possible number of fixed points for elements in the same isotopy class. Hence it is natural to study the image $[G]$ of a group G in certain mapping class groups. Anyway if we replace blindly any element of the group G with a simpler one in its isotopy class we lost the algebraic structure; the new group is in general not nilpotent even if G was. We need to realize $[G]$ as a nilpotent group of orientation-preserving homeomorphisms.

Definition 2.10. Let S be a compact surface, perhaps finitely punctured, with negative Euler characteristic and A be a subgroup of the mapping class group of S. We say that A is irreducible if every element of A is either pseudo-Anosov or of finite order.

Lemma 2.1. *Let S be a compact surface, perhaps finitely punctured, with negative Euler characteristic and A be an irreducible nilpotent subgroup of the mapping class group of S. Then there is a homomorphism $\sigma : A \to Homeo+(S)$ such that $\sigma(\theta)$ is a canonical Thurston representative in the mapping class θ. Moreover if A contains a pseudo-Anosov element then $\sigma(A)$ is generated by a finite group and a pseudo-Anosov element.*

Proof. The group A is solvable. Hence it is virtually abelian and finitely generated (Theorems A and B of [2]). If there is no pseudo-Anosov class in the center of A then A is finite (every finitely generated nilpotent group with finite center is finite, see Lemma 0.1, page 3 [1]). There exists a Riemannian metric in S and homomorphism σ from A to the isometries of S such that for each $\alpha \in A$ the element $\sigma(\alpha)$ is in the mapping class of α (Kerchoff's solution of the Nielsen realization problem [13]). The realization of the center of a pseudo-Anosov mapping class is solved in Lemma 2.10 of [9]. Indeed the center of a pseudo-Anosov class $[f_0]$ is generated by its subgroup of finite order elements and a pseudo-Anosov class $[f_1]$. □

Remark 2.2. In this paper we always apply Lemma 2.1 to finitely punctured spheres. In such a case we can describe the nature of the subgroup of finite order elements of $\sigma(A)$. The list of finite groups of orientation-preserving homeomorphisms defined in the sphere is well known. It coincides up to conjugacy with the list of finite groups of orientation preserving isometries. The list is composed by cyclic groups C_n, dihedral groups D_n with $2n$ elements, A_4, S_4 and S_5 (see page 40 [21]). The unique nilpotent groups in that list are the cyclic groups and the dihedral groups D_{2^n} for $n \in \mathbb{N}$.

Notice that if $\sigma(A)$ has non-trivial finite order elements so does the center of A. Since finite order elements have exactly two fixed points we deduce that $\sigma(A)$ has either a finite orbit of 2 elements or 2 fixed points. Moreover if $\sigma(A)$ has a global fixed point then it has a second one. If $\sigma(A)$ has no torsion elements then it is generated by a pseudo-Anosov element and it has at least 2 fixed points.

2.4. **Isotopies and fixed points.** The next results are key ingredients to translate results for simple models (provided by Lemma 2.1) to the initial nilpotent subgroup of $\mathrm{Diff}_+^1(\mathbb{S}^2)$.

Proposition 2.3. *(Proposition 4.2* [9]*) Suppose that* $\theta : S \to S$ *is an orientation-preserving homeomorphism of a finitely punctured surface, that* C *is a compact invariant set , that* M_i *is an essential connected* θ*-invariant subsurface of* $M := S \setminus C$ *with finite negative Euler characteristic and that* $\theta_{|M_i}$ *is either pseudo-Anosov relative to its puncture set or has finite order greater than one. Suppose further that* $z \in \mathrm{Fix}(\theta)$ *is contained in the interior of* $M_i \setminus C$ *and that* $f_t : S \to S$ *is an isotopy rel* C *from* $f_0 = \theta$ *to some* $f = f_1$*. Let* μ *be the Nielsen class for* θ *relative to* C *determined by* z *and let* ν *be the Nielsen class for* f *relative to* C *determined by* μ *and* f_t*. Then* ν *is non-trivial, compactly covered and does not peripherally contain any punctures.*

Suppose that $\mathcal{F}$ is a finitely generated group of orientation-preserving homeomorphisms defined in a finitely punctured sphere S. Consider a compact $\mathcal{F}$-invariant subset $K \subset S$ and a connected essential subsurface M_1 of $S \setminus K$ that is $\mathcal{F}$-invariant up to isotopy rel K and has negative Euler characteristic. Let M be the connected component of $S \setminus K$ containing M_1. There is a homeomorphism $\eta : \mathcal{F} \to \mathrm{MCG}(M_1)$ defined by $\phi \mapsto [f_{|M_1}]$ where f is a homeomorphism preserving M_1 that is isotopic to ϕ rel K.

Proposition 2.4. *Consider the notations above. Suppose that* $\mathcal{F}$ *is nilpotent and* $\eta(\mathcal{F})$ *is a non-trivial irreducible group. Let* Θ *be the model for* $\eta(\mathcal{F})$ *provided by Lemma 2.1. For each* $z \in \mathrm{Fix}(\Theta)$ *that is contained in the interior of* $M_1 \setminus K$ *there exists a lift* $\tilde{\mathcal{F}}$ *of* $\mathcal{F}$ *to the universal covering of* $M \setminus K$ *such that given a non-trivial* $\eta(\phi)$ *the set* $\mathrm{Fix}(\tilde{\phi})$ *is compact and non-empty.*

Suppose further that $\mathcal{F} =< Z^{(k)}(\mathcal{F}), \phi_1, \ldots, \phi_p >$ *is a nilpotent group of nilpotency class less or equal than* $k+1$*. Then there exist* α_1*, …,* α_p *in* $\mathcal{F}$ *such that* $\tilde{\mathcal{F}} =< Z^{(k)}(\tilde{\mathcal{F}}), \tilde{\alpha}_1, \ldots, \tilde{\alpha}_p >$ *and* $\mathrm{Fix}(\tilde{\alpha}_j)$ *is a non-empty compact set for any* $1 \leq j \leq p$*.*

Proof. Let $\tilde{M}_1$, $\tilde{M}$ be the universal coverings of M_1, M respectively. Fix a lift $\tilde{z}$ of z in $\tilde{M}_1$. There exists a unique lift $\tilde{\theta}$ of θ to $\tilde{M}_1$ such that $\tilde{\theta}(\tilde{z}) = \tilde{z}$ for any $\theta \in \Theta$. We obtain a lift $\tilde{\Theta}$ of Θ.

Given $\phi \in \mathcal{F}$ consider an element $\theta \in \mathrm{Homeo}_+(S)$ that is isotopic to ϕ rel K and satisfies $\theta_{|M_1} \in \Theta$. The restriction $\theta_{|M_1}$ is uniquely defined. By lifting the isotopy from θ to ϕ we obtain a lift $\tilde{\phi}_1 : \tilde{M}_1 \to \tilde{M}$ that extends uniquely to a homeomorphism $\tilde{\phi} : \tilde{M} \to \tilde{M}$. The uniqueness of the construction implies that $\phi \mapsto \tilde{\phi}$ defines a lift $\tilde{\mathcal{F}}$ of $\mathcal{F}$. Proposition 2.3 implies that $\mathrm{Fix}(\tilde{\phi})$ is a non-empty compact set if $\eta(\phi)$ is not a trivial class.

There exists $\beta \in \mathcal{F}$ such that $\eta(\beta)$ is non-trivial. Moreover we can suppose that β belongs to $Z^{(k)}(\mathcal{F}) \cup \{\phi_1, \ldots, \phi_p\}$. We define $\alpha_j = \phi_j$ if $\eta(\phi_j)$ is non-trivial and $\alpha_j = \phi_j \beta$ otherwise. $\square$

3. Fixed point theorem for the plane

Theorem 1.3 for commutative groups in $\mathrm{Diff}_{+}^{1}(\mathbb{R}^2)$ was proved in [9]. In this section we extend the result to nilpotent groups. More precisely, we prove the result for subgroups of $\mathrm{Diff}_{+}^{1}(\mathbb{R}^2)$ in $\mathcal{G}_{k,p}$ by a double induction process, first on p then on k.

Let us explain the idea of the proof. We consider a nilpotent subgroup G of $\mathrm{Diff}_{+}^{1}(\mathbb{R}^2)$ as a subgroup of $\mathrm{Homeo}_{+}(\mathbb{S}^2)$ where all elements fix the point at infinity. At this point we consider a G-invariant compact set K, for instance the fixed point set of a normal subgroup H of G. The idea is taking profit of a decomposition à la Thurston (with improvements along the lines in [9]) of the classes of isotopy rel K. Roughly speaking, the set of isolated points of K is finite and suitable to apply a Thurston decomposition whereas the set of accumulation points K' of K satisfies that any element of H is isotopic to the identity map in the neighborhood of K' rel K' (this property is a consequence of the smoothness of the elements of G and does not hold true for homeomorphisms). Hence it is natural to require K to be compact in $\mathbb{R}^2$ since the elements of G are not C^1 at ∞. A technical difficulty arises since in general the set $\mathrm{Fix}(H)$ is not compact. The following statements $A_{k,p}$, $B_{k,p}$, $C_{k,p}$ have hypotheses requiring the group to have varying degrees of compactness:

$\mathbf{A_{k,p}}$: Th. 1.3 holds true for any group $G = < Z^{(k)}(G), \phi_1, \ldots, \phi_p >$ of nilpotency class $\leq k+1$ if $\mathrm{Fix}(\phi_l)$ is compact for any $1 \leq l \leq p$.

$\mathbf{B_{k,p}}$: Th. 1.3 holds true for any group $G = < Z^{(k)}(G), \phi_1, \ldots, \phi_p >$ of nilpotency class $\leq k+1$ if $\mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_{p-1} >$ is compact.

$\mathbf{C_{k,p}}$: Th. 1.3 holds true for any group $G = < Z^{(k)}(G), \phi_1, \ldots, \phi_p >$ of nilpotency class $\leq k+1$.

We can consider the hypotheses in Statements $A_{k,p}$, $B_{k,p}$, $C_{k,p}$ as conditions of strong, weak and no compactness respectively. The validity of $C_{0,0}$ is obvious whereas $C_{0,1}$ is a consequence of Brouwer translation theorem. Moreover $C_{k,0}$ implies $C_{k,1}$ for $k > 0$ since $< Z^{(k)}(G), \phi >$ is a group of nilpotency class less or equal than k for any $\phi \in G$.

3.1. **Thurston normal form.** In this section we provide a Thurston decomposition for nilpotent groups of C^1 diffeomorphisms.

Proposition 3.1. *Suppose that S is a finitely punctured surface. Let G be a finitely generated nilpotent group of* $\mathrm{Diff}_{+}^{1}(S)$. *Consider normal* *subgroups* H_1*,* H_2 *of* G *and* G*-invariant compact sets* K_1*,* K_2 *such that* $K_1 \subset \mathrm{Fix}(H_1)$*,* $K_2 \subset \mathrm{Fix}(H_2) \setminus \mathrm{Fix}(H_1)$ *and* $K_1 \cap K_2 = \emptyset$*. Suppose that one of the following properties holds true:*

- K_2 *is finite.*
- $G \in \mathcal{G}_{k,p+1}$*,* $H_1 \in \mathcal{G}_{k,p}$*,* $K_1 = \mathrm{Fix}(H_1)$*,* $\mathrm{Fix}(H_1) \cap \mathrm{Fix}(H_2) = \emptyset$ *and* $C_{k,p}$ *holds true.*

Denote $M := S \setminus (K_1 \cup K_2)$*. Suppose that* $\chi(M) < 0$ *if* $\sharp(K_1 \cup K_2) < \infty$*.*

Then there is a finite set R *of disjoint simple closed curves (called reducing curves) in* M *and for any* $\phi \in G$ *there is a homeomorphism* $\theta : S \to S$ *isotopic to* ϕ *rel* $K_1 \cup K_2$ *such that*

1. θ *permutes disjoint open annulus neighborhoods in* S *of the elements of* R*.*

   Denote the union of the annular neighborhoods by $\mathcal{A}$*, let* $\{S_i\}$ *be the components of* $S \setminus \mathcal{A}$*, let* $X_i = (K_1 \cup K_2) \cap S_i$ *and let* $M_i := S_i \setminus X_i$ *. Then we have:*
2. *If* X_i *is infinite then either* $X_i \subset K_1$ *or* $X_i \subset K_2$*. In the former case* $\theta_{|S_i} \equiv Id$ *if* ϕ *belongs to* H_1*. In the latter case* $\theta_{|S_i} \equiv Id$ *if* ϕ *belongs to* H_2*.*
3. *If* X_i *is finite then* M_i *has negative Euler characteristic and* θ^{r_i} *is either periodic or pseudo-Anosov, where* r_i *is the smallest positive integer such that* $\theta^{r_i}(M_i) = M_i$*. Moreover, the* θ^{r_i} *for* $\phi \in G$ *generate a nilpotent subgroup that is either finite or virtually cyclic, meaning that it has a finite index cyclic subgroup.*

This result is the analogue of Proposition 2.11 of [9] adapted to the nilpotent setting. The proof is also similar. We sketch the proof, following the presentation of [9] and stressing the points where changes are required.

Definition 3.1. We define the dual tree of R. The vertices are the connected components of $S \setminus \mathcal{A}$. We define an edge for any $\gamma \in R$; it connects the two connected components of $S \setminus \mathcal{A}$ that share boundary components with the annulus neighborhood of γ. If we assign length one to every edge, the group G acts on the tree by isometries.

Suppose that S is a finitely punctured surface. Assume that $\mathcal{F}$ is a finitely generated subgroup of $\mathrm{Diff}^1_+(S)$ and B, C are $\mathcal{F}$-invariant compact subsets of F such that $B \subset \mathrm{Fix}(\mathcal{F})$, $C \cap \mathrm{Fix}(\mathcal{F}) = \emptyset$. Suppose that the image of $\mathcal{F}$ in the group of classes of homeomorphisms modulo isotopy rel $B \cup C$ is nilpotent.

Definition 3.2. Let $\mathcal{W}(\mathcal{F}, B, C)$ be the set of compact subsurfaces (they can contain punctures) $W \subset S$ such that:

- $(\mathcal{W}-1)$ ∂W has finitely many components, each of which is contained in and is essential in $M := S \setminus (B \cup C)$;
- $(\mathcal{W}-2)$ W contains all but finitely many points of B and every component of W intersects B in an infinite set;
- $(\mathcal{W}-3)$ for all $f \in \mathcal{F}$ the following is satisfied:
  - $(\mathcal{W}-3f)$ there exists $\phi : S \to S$ isotopic to f rel $B \cup C$ such that $\phi_{|W}$ is the identity.

The set $\mathcal{W}(\mathcal{F}, B, C)$ is not empty if B is an infinite set. This is Lemma 2.5 of [9] for the commutative case. The proof holds true for any finitely generated subgroup of $\mathrm{Diff}_+^1(S)$.

Consider the setup in Proposition 3.1. The idea is that a maximal element W of $\mathcal{W}(\mathcal{F}, B, C)$ is unique up to isotopy. If we define $\mathcal{F} = H_1$, $B = K_1$ and $C = K_2$ the uniqueness implies that W is invariant by G up to isotopy rel $K_1 \cup K_2$. Naturally the connected components of W are pieces of the decomposition, they satisfy the condition (2) in Proposition 3.1.

Let us prove that $\mathcal{W}(\mathcal{F}, B, C)$ has maximal elements. A partial order can be defined in $\mathcal{W}(\mathcal{F}, B, C)$. We have $W_1 < W_2$ for $W_1, W_2 \in \mathcal{W}(\mathcal{F}, B, C)$ if W_1 is isotopic rel $B \cup C$ to a subsurface of W_2 but W_1 is not isotopic rel $B \cup C$ to W_2. The next lemma is a version of Lemma 2.6 of [9] adapted to the nilpotent setting.

Lemma 3.1. *Suppose that B is infinite and that either of the following is satisfied.*

- *C is finite.*
- *$\mathcal{F}$ belongs to $\mathcal{G}_{k,p}$, $C_{k,p}$ holds true and $B = \mathrm{Fix}(\mathcal{F})$.*

Then $\mathcal{W}(\mathcal{F}, B, C)$ has maximal elements.

Proof. The proof for the case $\sharp C < \infty$ is the same as in Lemma 2.6 of [9]. Moreover they prove that if the result is false there exists a disc D in S that is $\mathcal{F}$-invariant modulo isotopy rel $B \cup C$, disjoint from B and such that $C_D := C \cap D$ is non-empty.

Let X be the universal covering of $S \setminus (B \cup (C \setminus C_D))$. We choose a lift $\tilde{D}$ of D to X. It is also a disc. Given $f \in \mathcal{F}$ there exists a unique lift $\tilde{f}$ of f to X such that $\tilde{f}(\tilde{C}_D) = \tilde{C}_D$ where $\tilde{C}_D$ is the lift of C_D to $\tilde{D}$. In this way we obtain a lift $\tilde{\mathcal{F}}$ of $\mathcal{F}$ to X that preserves the compact set $\tilde{C}_D$. Since $C_{k,p}$ holds true then $\mathrm{Fix}(\tilde{\mathcal{F}})$ is non-empty. Therefore $\mathrm{Fix}(\mathcal{F}) \cap (S \setminus B)$ is not empty. This is absurd. □

The Thurston decomposition for nilpotent groups is proved as a part of the inductive step (we need to use that $C_{k,p}$ holds true). This is not necessary in [9] since in the case $\sharp C = \infty$ the group $\mathcal{F}$ is always cyclic and the contradiction is provided by the Brouwer translation theorem.

The set $\mathcal{W}(\mathcal{F}, B, C)$ has a unique maximal element, i.e. two maximal elements of $\mathcal{W}(\mathcal{F}, B, C)$ are isotopic rel $B \cup C$. The proofs in Lemmas 2.7 and 2.8 of [9] apply here.

Proof of Proposition 3.1. We follow the steps of Proposition 2.11 of [9]. We include a proof for the sake of clarity.

We define $W_1 = \emptyset$ if K_1 is finite, otherwise consider a maximal element W_1 of $\mathcal{W}(H_1, K_1, K_2)$. Since H_1 is a normal subgroup of G and K_1, K_2 are G-invariant we obtain that $\phi(W_1)$ is isotopic to W_1 rel $K_1 \cup K_2$ for any $\phi \in G$. There exists a diffeomorphism $\theta_\phi : S \to S$ such that ϕ is isotopic to θ_ϕ rel $K_1 \cup K_2$ and $\theta_\phi(W_1) = W_1$ for any $\phi \in G$. Moreover we can suppose that $(\theta_\phi)_{|W_1} \equiv Id$ if $\phi \in H_1$.

We denote the finitely puncture surface $S \setminus W_1$ by S'. Consider a set of generators $\{\phi_1, \ldots, \phi_m\}$ of G such that $H_2 =< \phi_1, \ldots, \phi_n >$ for some $n \leq m$. We can suppose that $\{\phi_1, \ldots, \phi_m\}$ and $\{\phi_1, \ldots, \phi_n\}$ contain set of generators of H_1 and $H_1 \cap H_2$ respectively. We denote $G' =< \theta_{\phi_1}, \ldots, \theta_{\phi_m} >$ and $H_2' =< \theta_{\phi_1}, \ldots, \theta_{\phi_n} >$. They are subgroups of $\mathrm{Diff}_+^1(S')$. Moreover H_2' is a normal subgroup of G' modulo isotopy rel $S' \cap (K_1 \cup K_2)$. If K_2 is finite we define $W_2 = \emptyset$. Otherwise consider a maximal element W_2 of $\mathcal{W}(H_2', K_2, K_1 \setminus W_1)$. We can suppose that $(\theta_{\phi_j})_{|W_2} \equiv Id$ for any $1 \leq j \leq n$ up to an isotopy rel $K_1 \cup K_2$. The maximal nature of W_2 implies that up to an isotopy rel $K_1 \cup K_2$ the diffeomorphism θ_{ϕ_j} preserves W_2 for any $1 \leq j \leq m$.

A partial set R' of reducing curves is obtained by removing from $\partial W_1 \cup \partial W_2$ the curves bounding an unpunctured annulus and replacing them with a core curve of the annulus. In this way we can suppose that G satisfies (1) and (2) with respect to R'. We need to find a set of reducing curves R containing R' such that (1) still is satisfied and (3) holds true. We consider the finitely puncture subsurface $S'' := S \setminus (W_1 \cup W_2)$ of S. The image G_0 of G in the mapping class group of S'' is a nilpotent group. In particular it is virtually abelian (Theorem B of [2]). An element of the mapping class group is irreducible if and only if an iterate is irreducible. Hence it suffices to prove the existence of a Thurston normal form for an abelian group of the mapping class group of a finitely punctured surface. Such result is well known, see for example Lemma 2.2 of [12]. The mapping class group defined by the stabilizer of every finite type subsurface of the decomposition is either finite or virtually cyclic by Lemma 2.1. □

3.2. **Proof of Theorem 1.3.** Our point of view is replacing a subgroup G of $\mathrm{Diff}_+^1(\mathbb{R}^2)$ with an irreducible nilpotent model that defines the same image in the mapping class group of a subsurface of $\mathbb{R}^2$, obtaining information for the model and then interpreting it for the initial group. One of the reasons because this approach works is that nilpotent groups are quite rigid. For instance the next lemma implies that in Theorem 1.3 we can replace the condition of existence of a G-invariant non-empty compact set with the existence of an element of G whose fixed point set is a non-empty compact set. A condition on a single element implies the existence of a global fixed point for the group.

Lemma 3.2. *Let* $G =< Z^{(k)}(G), \phi_1, \ldots, \phi_p >$ *be a finitely generated nilpotent subgroup of* $\mathrm{Diff}_+^1(\mathbb{R}^2)$ *of nilpotency class* $\leq k+1$. *Suppose that there exists* $\alpha \in G$ *such that* $\mathrm{Fix}(\alpha)$ *is a non-empty compact set. Suppose further that* $C_{k,p}$ *holds true. Then* $\mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_p >$ *is non-empty.*

Proof. The series

$$1 \leq < Z^{(0)}(G), \alpha > \leq < Z^{(1)}(G), \alpha > \leq \ldots \leq < Z^{(k)}(G), \alpha >$$

is subnormal and every quotient of consecutive subgroups is abelian. Let us prove that $\mathrm{Fix} < Z^{(l)}(G), \alpha >$ is non-empty for any $0 \leq l \leq k$. It is obvious for $l = 0$ and if $l < k$ the compact set $\mathrm{Fix} < Z^{(l)}(G), \alpha >$ is invariant by $< Z^{(l+1)}(G), \alpha >$. We obtain $\mathrm{Fix} < Z^{(l+1)}(G), \alpha > \neq \emptyset$ by Statement $C_{k,0}$. Since $< Z^{(k)}(G), \alpha >$ is normal in G, Statement $C_{k,p}$ implies that $\mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_p >$ is not empty. □

Let $G =< Z^{(k)}(G), \phi_1, \ldots, \phi_{p+1} >$ in $\mathcal{G}_{k,p+1}$. Consider $a_1 < a_2 < \ldots < a_{k+1}$ and a system of generators $\{g_1, \ldots, g_{a_{k+1}}\}$ of G such that $Z^{(l)}(G) =< g_1, \ldots, g_{a_l} >$ for any $1 \leq l \leq k+1$, $a_{k+1} - a_k = p+1$ and $g_{a_k+1} = \phi_1$, …, $g_{a_{k+1}} = \phi_{p+1}$. We denote $G_0 = \{Id\}$ and $G_j =< g_1, \ldots, g_j >$. The group G_j is normal in G for any $0 \leq j < a_{k+1}$. We define $J_j = G_j$ for $j < a_{k+1} - 1$ and $J_{a_{k+1}-1} = G_{a_{k+1}-2}$ for $j = a_{k+1} - 1$.

Lemma 3.3. *Let* $K \subset \mathrm{Fix}(G_j)$ *be a non-empty compact* G*-invariant subset for* $j < a_{k+1}$. *Suppose that* $C_{k,p}$ *holds true. Then one of the following possibilities is satisfied:*

1. $K \cap \mathrm{Fix}(G_{j+1}) \neq \emptyset$.
2. *There exists a lift* $\tilde{G}$ *of* G *such that* $\tilde{G} =< Z^{(k)}(\tilde{G}), \tilde{\alpha}_1, \ldots, \tilde{\alpha}_{p+1} >$ *and* $\mathrm{Fix}(\tilde{\alpha}_l)$ *is a non-empty compact set for any* $1 \leq l \leq p+1$.
3. $\cap_{\phi \in G} \phi(P(K, g_{j+1}) \cap \mathrm{Fix}(J_j))$ *is a non-empty compact* G*-invariant set. Moreover if* $j = a_{k+1} - 1$ *then there exists a lift* $\tilde{G}$ *of* G *such that* $\mathrm{Fix}(\tilde{\phi}_{p+1})$ *is non-empty and compact.*

This lemma justifies the introduction of Statements $A_{k,p}$ and $B_{k,p}$. In Cases (1) and (3) for $j < a_{k+1} - 1$ there exists a compact non-empty G-invariant set F contained in $\mathrm{Fix}(G_{j+1})$. We can continue the process by replacing K with F and G_j with G_{j+1}. Otherwise either (2) holds true and $\tilde{G}$ satisfies the hypothesis in Statement $A_{k,p+1}$ or (3) holds true for $j = a_{k+1} - 1$ and $\tilde{G}$ satisfies the hypothesis in Statement $B_{k,p+1}$.

Proof. We can suppose that $K \cap \mathrm{Fix}(G_{j+1}) = \emptyset$. Consider a compact set $K^1 \subset K$ that is minimal by the action of G. We apply Prop. 3.1 to $H_1 = G_j$, $H_2 = G$, $K_1 = K^1$, $K_2 = \emptyset$. Consider the tree Γ that is dual to R (see Def. 3.1). Then G fixes the vertex that corresponds to the connected component D_∞ of ∞. Consider the subtree Γ' of Γ of points fixed by G. If $\Gamma' = \{D_\infty\}$ we define $D = D_\infty$. Otherwise there exists a vertex $D \neq D_\infty$ of Γ' that belongs to exactly one edge γ of Γ'.

Suppose that $D \cap K^1$ is finite. Let $\hat{D}$ be the marked sphere obtained from $\overline{D}$ by contracting $\overline{D} \cap \mathcal{A}$. Consider $[] : G \to \mathrm{MCG}(G_{|\hat{D}})$. The model for $[G]$ provided by Lemma 2.1 has exactly a fixed point at $\overline{D} \cap (\mathcal{A} \cup \{\infty\})$ and it does not fix any other puncture or curve in $\mathcal{A}$. Hence it fixes an interior point (Remark 2.2). There exists a lift $\tilde{G}$ of G to the universal covering of the connected component of $\mathbb{R}^2 \setminus K^1$ containing $D \setminus K^1$ such that $\tilde{G} =< Z^{(k)}(\tilde{G}), \tilde{\alpha}_1, \ldots, \tilde{\alpha}_{p+1} >$ and $\mathrm{Fix}(\tilde{\alpha}_l)$ is a non-empty compact set for any $1 \leq l \leq p+1$ (Prop. 2.4).

Suppose $D \cap K^1$ is an infinite set. The minimality implies $K^1 \subset D$ and $R = \emptyset$. In particular all the elements in G_j are isotopic to the identity rel $K^1 \cup \{\infty\}$.

Consider a compact set $K^2 \subset K^1$ that is minimal by the action of G_{j+1}. Let E_∞ be the unbounded connected component of $\mathbb{R}^2 \setminus K^2$. Consider the full compact set $\hat{K}^2 := \mathbb{R}^2 \setminus E_\infty$. Suppose that $\mathrm{Fix}(g_{j+1}) \cap \hat{K}^2$ is non-empty. Then there exists a bounded connected component D_1 of $\mathbb{R}^2 \setminus K^2$ such that $\mathrm{Fix}(g_{j+1}) \cap D_1 \neq \emptyset$. Moreover the minimality implies $K^2 \subset \overline{E_\infty}$, hence D_1 is a topological disc. The diffeomorphism g_{j+1} has no fixed points in ∂D_1; thus the set $\mathrm{Fix}((g_{j+1})_{|D_1})$ is compact. Since D_1 is G_{j+1}-invariant we obtain that $\mathrm{Fix} < J_j, g_{j+1} > \cap D_1$ is non-empty by Lemma 3.2. Clearly $\mathrm{Fix} < J_j, g_{j+1} > \cap D_1$ and all its iterates by elements of G are contained in $P(K^1, g_{j+1}) \cap \mathrm{Fix}(J_j)$. Moreover we define $\tilde{G}_{j+1}$ as the group obtained by restricting G_{j+1} to D_1. We suppose $\mathrm{Fix}(g_{j+1}) \cap \hat{K}^2 = \emptyset$ from now on. The set $P(K^2, g_{j+1})$ is equal to $P(\hat{K}^2, g_{j+1})$, it is a non-empty subset of E_∞ (Prop. 2.2). Since $\hat{K}^2$ is a full compact set then it is not connected, otherwise we obtain $P(\hat{K}^2, g_{j+1}) = \emptyset$. In particular K_2 is not connected.

Let $\sigma : \tilde{E}_\infty \to E_\infty$ be the universal covering of E_∞. Given $\phi \in G_j$ let $\{\beta_t\}_{t \in [0,1]}$ be an isotopy rel K^2 from $\beta_0 = Id$ to $\beta_1 = \phi$. Consider the lift $\{\tilde{\beta}_t\}_{t \in [0,1]}$ to $\tilde{E}_\infty$ such that $\tilde{\beta}_0 = Id$. We define $\tilde{\phi} = \tilde{\beta}_1$. It is the identity lift of $\phi \in G_j$ to $\tilde{E}_\infty$; it is uniquely defined since the Euler characteristic of E_∞ is negative. We define $\mathrm{Fix}_c(\phi) = \sigma(\mathrm{Fix}(\tilde{\phi}))$ and $\mathrm{Fix}_c(J_j) = \cap_{\phi \in J_j} \mathrm{Fix}_c(\phi)$. Let us prove that $P(K^2, g_{j+1}) \cap \mathrm{Fix}_c(J_j)$ is contained in $\phi(P(K, g_{j+1}) \cap \mathrm{Fix}(J_j))$ for any $\phi \in G$. Given $\phi \in G$ we have

$$\alpha := \phi g_{j+1} \phi^{-1} g_{j+1}^{-1} \in G_j \cap Z^{(k)}(G) \subset J_j.$$

We claim

$$P(K^2, \alpha g_{j+1}) \cap \mathrm{Fix}_c(J_j) = P(K^2, g_{j+1}) \cap \mathrm{Fix}_c(J_j). \tag{1}$$

Indeed if an element z of $\mathrm{Fix}_c(J_j) \cap \mathrm{Fix}(g_{j+1})$ does not belong to $P(K^2, \alpha g_{j+1})$ then there exists a path ρ from z to ∞ in $\mathbb{S}^2 \setminus K^2$ such that ρ and $(\alpha g_{j+1})(\rho)$ are homotopic relative to ends in $\mathbb{S}^2 \setminus K^2$. Let $\{\alpha_t\}_{t \in [0,1]}$ be an isotopy rel $K^1 \cup \{\infty\}$ such that $\alpha_0 = Id$ and $\alpha_1 = \alpha$. We obtain that ρ and the path $\varsigma g_{j+1}(\rho)$ are homotopic relative to their ends in $\mathbb{S}^2 \setminus K^2$ where $\varsigma : [0,1] \to \mathbb{R}^2 \setminus K^1$ is defined by $\varsigma(t) = \alpha_t^{-1}(z)$. The class of ς is trivial in $\pi_1(\mathbb{R}^2 \setminus K^2, z)$ since $z \in \mathrm{Fix}_c(\alpha)$. Thus z does not belong to $P(K^2, g_{j+1})$. Analogously we prove that $z \in (\mathrm{Fix}_c(J_j) \cap \mathrm{Fix}(g_{j+1})) \setminus P(K^2, g_{j+1})$ implies $z \notin P(K^2, \alpha g_{j+1})$.

We deduce

$$P(K^2, g_{j+1}) \cap \mathrm{Fix}_c(J_j) \subset P(\phi(K), \alpha g_{j+1}) \cap \mathrm{Fix}(J_j) =$$

$$P(\phi(K), \phi g_{j+1} \phi^{-1}) \cap \mathrm{Fix}(J_j) = \phi(P(K, g_{j+1}) \cap \mathrm{Fix}(J_j)).$$

The inclusion is a consequence of equation (1) and $K^2 \subset K$. In order to prove that $\cap_{\phi \in G} \phi(P(K, g_{j+1}) \cap \mathrm{Fix}(J_j))$ is non-empty it suffices to show that $P(K^2, g_{j+1}) \cap \mathrm{Fix}_c(J_j)$ is non-empty.

We apply the Thurston decomposition (Prop. 3.1) to the group G_{j+1} and the marked surface $(\mathbb{S}^2, \{\infty\})$ ($H_1 = G_j$, $H_2 = G_{j+1}$, $K_1 = K^2$, $K_2 = \emptyset$). Let D'_∞ be the connected component of $\mathbb{S}^2 \setminus \mathcal{A}$ that contains ∞. Suppose that $D'_\infty \cap K^2$ is of finite type. The minimality implies that g_{j+1} can not fix any curve in $\overline{D'_\infty} \cap \mathcal{A}$. Analogously as in the case $\sharp(D \cap K^1) < \infty$ Prop. 2.4 implies the existence of a lift $\tilde{G}_{j+1}$ of G_{j+1} to $\tilde{E}_\infty$ such that $\mathrm{Fix}(\tilde{g}_{j+1})$ is a non-empty compact set. Moreover $\sigma(\mathrm{Fix}(\tilde{g}_{j+1}))$ is contained in $P(K^2, g_{j+1})$. We obtain $\mathrm{Fix} < \tilde{J}_j, \tilde{g}_{j+1} > \neq \emptyset$ by Lemma 3.3. The set $P(K^2, g_{j+1}) \cap \mathrm{Fix}_c(J_j)$ is not empty, it contains $\sigma(\mathrm{Fix} < \tilde{J}_j, \tilde{g}_{j+1} >)$.

Suppose that $D'_\infty \cap K^2$ is not of finite type. In this case K^2 is contained in D'_∞ by minimality and R is empty. If there is an essential finite type subsurface $N \subset \mathbb{R}^2 \setminus K^2$ preserved by g_{j+1} modulo isotopy rel K^2 we can redefine R to be ∂N. The connected component D'_∞ of infinity in $\mathbb{S}^2 \setminus R$ satisfies $K^2 \cap D'_\infty = \emptyset$ by minimality. We can argue as in the finite type case.

Consider a lift $\tilde{z} \in \tilde{E}_\infty$ of a point z in $P(K^2, g_{j+1})$. Let $\tilde{g}_{j+1}$ be the lift of g_{j+1} to $\tilde{E}_\infty$ such that $\tilde{g}_{j+1}(\tilde{z}) = \tilde{z}$. Since g_{j+1} does not preserve an essential finite type subsurface, Lemma 2.12 of [9] implies that g_{j+1} does not fix the homotopy class of any essential non-peripheral closed curve. It follows that $\mathrm{Fix}(\tilde{g}_{j+1})$ is compact. No iterate of g_{j+1} belongs to G_j, otherwise K^2 is finite. The mapping $\tau : G_{j+1} \to < g_{j+1} >$ defined by $\tau_{|G_j} \equiv Id$, $\tau(g_{j+1}) = g_{j+1}$ is a well-defined homomorphism of groups. All the elements of G_j are isotopic to the identity rel K^2. Given $\phi \in G_{j+1}$ let $\{\beta_t\}_{t \in [0,1]}$ be an isotopy rel K^2 from $\beta_0 = \tau(\phi)$ to $\beta_1 = \phi$. Consider the lift $\{\tilde{\beta}_t\}_{t \in [0,1]}$ to $\tilde{E}_\infty$ such that $\tilde{\beta}_0(\tilde{z}) = \tilde{z}$. We define $\tilde{\phi} = \tilde{\beta}_1$. The group $\tilde{G}_{j+1}$ is a lift of G_{j+1} to $\tilde{E}_\infty$ by the uniqueness of the construction. Moreover $\tilde{G}_j$ is the identity lift of G_j. Lemma 3.2 implies $\mathrm{Fix} < \tilde{J}_j, \tilde{g}_{j+1} > \neq \emptyset$. The Nielsen class of the elements of $\sigma(\mathrm{Fix}(\tilde{g}_{j+1}))$ is the Nielsen class of z. Thus $P(K^2, g_{j+1}) \cap \mathrm{Fix}_c(J_j)$ is non-empty. $\square$

Let us prove next $C_{k,p} \implies A_{k,p+1}$ and $C_{k,p} \implies B_{k,p+1}$. The proof of $C_{k,p} \implies C_{k,p+1}$ is an easy consequence of these results and Lemma 3.3.

Lemma 3.4. *Statement $C_{k,p}$ implies Statement $A_{k,p+1}$ for all $k \geq 0$, $p \geq 0$.*

Proof. Denote

$$H_1 =< Z^{(k)}(G), \phi_1, \ldots, \phi_p >, \quad H_2 =< Z^{(k)}(G), \phi_1, \ldots, \phi_{p-1}, \phi_{p+1} > .$$

We define $K_1 = \mathrm{Fix}(H_1)$ and $K_2 = \mathrm{Fix}(H_2)$.

The hypothesis implies that K_1 and K_2 are compact. Moreover $C_{k,p}$ implies that none of them is empty. It suffices to prove that $K_1 \cap K_2 = \emptyset$ leads to a contradiction.

We apply Prop. 3.1 to H_1, H_2. Consider the tree Γ that is dual to R. We define D as in Lemma 3.3. It is a G-invariant (modulo isotopy) component of $\mathbb{S}^2 \setminus \mathcal{A}$ such that there is exactly one point or curve γ fixed by G in the union of $\{\infty\} \cup (\overline{D} \cap \mathcal{A})$. If $D = D_\infty$ we define $D' = \mathbb{R}^2$. Otherwise we define D' as the connected component of $\mathbb{S}^2 \setminus \gamma$ not containing ∞. We have $D \subset D'$.

Suppose $K_1 \cap D' \neq \emptyset$ and $K_2 \cap D' \neq \emptyset$. If $\sharp(K_1 \cap D) = \infty$ then the elements in H_1 fix the curves in $\overline{D} \cap \mathcal{A}$. Since $\emptyset \neq K_2 \cap D' \subset \mathrm{Fix}(H_2)$ there exists a curve in $\overline{D} \cap \mathcal{A}$ different than γ and invariant by the group $G =< H_1, H_2 >$. This is impossible by the choice of D. Hence $\sharp(K_1 \cap D) = \infty$ implies $K_2 \cap D' = \emptyset$. Analogously $\sharp(K_2 \cap D) = \infty$ implies $K_1 \cap D' = \emptyset$. Either $(K_1 \cup K_2) \cap D$ is finite or one of the sets $K_1 \cap D'$, $K_2 \cap D'$ is empty.

Suppose that $(K_1 \cup K_2) \cap D$ is a finite set. Let $\hat{D}$ be the marked sphere obtained from $\overline{D}$ by contracting the curves in $\overline{D} \cap \mathcal{A}$. Consider $[] : G \to \mathrm{MCG}(G_{|\hat{D}})$. The model for $[G]$ provided by Lemma 2.1 has exactly a fixed point at $\overline{D} \cap (\mathcal{A} \cup \{\infty\})$ and it does not fix any other puncture or curve in $\mathcal{A}$. Hence it fixes an interior point (Remark 2.2). Proposition 2.4 implies that there exists a lift $\tilde{G}$ of G to the universal covering of the connected component of $\mathbb{R}^2 \setminus (K_1 \cup K_2)$ containing $D \setminus (K_1 \cup K_2)$ such that $\tilde{G} =< Z^{(k)}(\tilde{G}), \tilde{\alpha}_1, \ldots, \tilde{\alpha}_{p+1} >$ and $\mathrm{Fix}(\tilde{\alpha}_j)$ is a non-empty compact set for any $1 \leq j \leq p+1$. Lemma 3.2 implies that $\mathrm{Fix} < Z^{(k)}(\tilde{G}), \tilde{\alpha}_1 >$ is a non-empty $\tilde{G}$-invariant compact set. We obtain $\mathrm{Fix} < Z^{(k)}(\tilde{G}), \tilde{\phi}_1, \ldots, \tilde{\phi}_p > \neq \emptyset$ by Statement $C_{k,p}$. We have

$$\emptyset \neq \mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_p > \cap (\mathbb{R}^2 \setminus (K_1 \cup K_2)) = \emptyset.$$

It represents a contradiction.

Suppose $\sharp((K_1 \cup K_2) \cap D) = \infty$. Then either $\sharp(K_1 \cap D) = \infty$ and $K_2 \cap D' = \emptyset$ or $\sharp(K_2 \cap D) = \infty$ and $K_1 \cap D' = \emptyset$. Suppose that we are in the latter case without lack of generality. Clearly we have $D \neq D_\infty$. Consider a lift $\tilde{D}'$ of the disc D' to the universal covering M of the connected component of $\mathbb{R}^2 \setminus ((K_1 \cup K_2) \setminus D')$ containing D'. Let $\tilde{K}$ be the lift of $(K_1 \cup K_2) \cap D'$ in $\tilde{D}'$. There exists a unique lift $\tilde{G}$ of G to M such that $\tilde{\phi}(\tilde{K}) = \tilde{K}$ for any $\phi \in G$. The group $\tilde{G}$ preserves the compact set $\tilde{K}$; we obtain $\mathrm{Fix}(\tilde{H}_1) \neq \emptyset$ and $\mathrm{Fix}(\tilde{H}_2) \neq \emptyset$ by Statement $C_{k,p}$. We deduce that the intersection of $\mathrm{Fix}(H_1)$ and $\mathbb{R}^2 \setminus K_1$ is not empty; this is absurd. ☐

Lemma 3.5. *Statement $C_{k,p}$ implies Statement $B_{k,p+1}$ for all $k \geq 0$, $p \geq 0$.*

Proof. We define H_1, H_2 and K_1 as in Lemma 3.4. The set K_1 is compact by hypothesis. We apply Lemma 3.3. If $K_1 \cap \mathrm{Fix}(G)$ is not empty there is nothing to prove. If the second Case is satisfied then Lemma 3.4 implies $\mathrm{Fix}(G) \neq \emptyset$. So we can suppose that

$$K_2 := \cap_{\phi \in G} \phi(P(K_1, \phi_{p+1}) \cap \mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_{p-1} >)$$

is non-empty. We apply Prop. 3.1 to H_1, H_2.

The remainder of the proof is exactly the same as in Lemma 3.4 except for the case when $\sharp(K_1 \cap D) = \infty$ and $K_2 \cap D' = \emptyset$. We apply Lemma 3.3 to the compact G-invariant set $K_1 \cap D'$. If we are in the second Case we obtain $\mathrm{Fix}(G) \neq \emptyset$ by applying Lemma 3.4. We can suppose that

$$K_2' := \cap_{\phi \in G} \phi \left(P(K_1 \cap D', \phi_{p+1}) \cap \mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_{p-1} > \right)$$

is a non-empty compact G-invariant set contained in K_2. We obtain

$$K_2' = (K_2' \cap D') \cup (K_2' \setminus \overline{D'}) = K_2' \setminus \overline{D'} \subset \mathbb{R}^2 \setminus P(K_1 \cap D', \phi_{p+1})$$

since $\partial D'$ is invariant by ϕ_{p+1} modulo an isotopy rel $K_1 \cup K_2$. This contradicts the property $K_2' \subset P(K_1 \cap D', \phi_{p+1})$. □

Lemma 3.6. *Statement $C_{k,p}$ implies Statement $C_{k,p+1}$ for all $k \geq 0$, $p \geq 0$.*

Proof. Consider the notations above Lemma 3.3. Let K be a compact non-empty G-invariant set. We define

$$j = \max\{l \in \{0, \ldots, a_{k+1}\} : K \cap \mathrm{Fix}(G_j) \neq \emptyset\}.$$

If $j = a_{k+1}$ we are done. We apply Lemma 3.3 to $K \cap \mathrm{Fix}(G_j)$. The first Case is impossible by the choice of j. If the second Case is satisfied then $\mathrm{Fix}(G)$ is non-empty by Lemma 3.4. So we can suppose that we are in the third Case. If $j < a_{k+1} - 1$ then the set $\cap_{\phi \in G} \phi(P(K, g_{j+1}) \cap \mathrm{Fix}(G_{j+1}))$ is a non-empty compact G-invariant set contained in $\mathrm{Fix}(G_{j+1})$. Up to repeat the argument a finite number of times we obtain a compact non-empty compact G-invariant set K_1 contained in $\mathrm{Fix} < Z^{(k)}(G), \phi_1, \ldots, \phi_p >$. We apply Lemma 3.3 to K_1. Either G has a global fixed point or the last Case is satisfied. In the latter situation there exists a lift $\tilde{G}$ of G such that $\mathrm{Fix}(\tilde{\phi}_{p+1})$ is compact. Lemma 3.5 implies that $\tilde{G}$ (and then G) has a global fixed point. □

Remark 3.1. Notice that since we already proved Theorem 1.3 the condition requiring $C_{k,p}$ to be satisfied in Proposition 3.1 and Lemmas 3.1, 3.2 and 3.3 is superfluous.

Proof of Corollary 1.1. By a theorem of Margulis a subgroup $\mathcal{F}$ of $\mathrm{Homeo}_+(\mathbb{S}^1)$ either contains a non abelian free subgroup or there exists a $\mathcal{F}$-invariant probability measure μ [17]. Consider the nilpotent group $\mathcal{F}$ of restrictions to $\mathbb{S}^1 = \partial \mathbb{D}^2$ of elements of G. Given an element $f \in \mathcal{F}$ such that $\mathrm{Fix}(f) \neq \emptyset$ the dynamics of f in each interval of $\mathbb{S}^1 \setminus \mathrm{Fix}(f)$ is conjugate to a translation. We obtain $\mathrm{supp}(\mu) \subset \mathrm{Fix}(f)$. Thus either $\mathrm{supp}(\mu) \subset \mathrm{Fix}(\mathcal{F})$ and there is nothing to prove or there exists $\phi \in G$ such that $\mathrm{Fix}(\phi) \cap \partial \mathbb{D}^2 = \emptyset$. In the latter case $\mathrm{Fix}(\phi)$ is compact in $\mathbb{D}^2 \setminus \partial \mathbb{D}^2$. Every finitely generated subgroup of G containing ϕ has a global fixed point in $\mathbb{D}^2 \setminus \partial \mathbb{D}^2$ by Lemma 3.2. Thus G has a global fixed point. □

Proposition 3.2. *Consider the notations in Prop. 2.4. Suppose that $\mathcal{F}$ is nilpotent and $\eta(\mathcal{F})$ is a non-trivial irreducible group. Let Θ be the model for $\eta(\mathcal{F})$ provided by Lemma 2.1. For each $z \in \mathrm{Fix}(\Theta)$ that* *is contained in the interior of* M_1 *there exists* $x \in \mathrm{Fix}(\mathcal{F}) \cap M$ *such that: given a non-trivial* $\eta(\phi)$ *the* ϕ*-Nielsen class rel* K *that contains* x *is compactly covered, does not peripherally contains any punctures and corresponds, via the isotopy between* θ_ϕ *and* ϕ*, to the* θ_ϕ*-Nielsen class rel* K *that contains* z *where* $\theta_\phi \in Homeo_+(S)$ *is isotopic to* ϕ *and* $(\theta_\phi)_{|M_1} \in \Theta$*.*

Proof. Proposition 2.4 provides a lift $\tilde{\mathcal{F}}$ of $\mathcal{F}$ to the universal covering $\sigma : \tilde{M} \to M$ of M. Suppose that $\eta(\phi)$ is not trivial, then $\mathrm{Fix}(\tilde{\phi})$ is a non-empty compact set. Moreover $\sigma(\mathrm{Fix}(\tilde{\phi}))$ corresponds to the θ_ϕ-Nielsen class rel K that contains z. Such a Nielsen class does not peripherally contain any punctures (Prop. 2.3).

It suffices to prove that $\mathrm{Fix}(\tilde{\mathcal{F}})$ is non-empty. This is a consequence of Lemma 3.2 and Theorem 1.3. □

4. Existence of orbits of period 2 in the sphere

Let us discuss briefly the proof of Theorem 1.2. It suffices to show the result for finitely generated nilpotent groups since a family of compact sets satisfying the finite intersection property has non-empty intersection. We will see that if Theorem 1.2 holds true for groups in $\mathcal{G}_{k,p}$ then any group $G \in \mathcal{G}_{k,p+1}$ has a finite orbit. The main ingredient of this result is Lemma 4.1 below. Once the finite G-orbit $\mathcal{O}$ is fixed we consider the Thurston decomposition of G in the marked surface $(\mathbb{S}^2, \mathcal{O})$. Roughly speaking we find an invariant connected component M of the decomposition and global fixed points for an irreducible nilpotent model of a normal subgroup of index at most 2 of G (see Remark 2.2). Then we lift this result to the initial group by using Proposition 2.4 and Theorem 1.3.

The following lemma is the generalization of Lemma 7.2 of [9] to the nilpotent case.

Lemma 4.1. *Suppose that* G *is a finitely generated nilpotent subgroup of* $\mathrm{Diff}^1_+(\mathbb{S}^2)$*, that* $X \subset \mathbb{S}^2$ *is a compact* G*-invariant set and that* $\gamma \subset M := \mathbb{S}^2 \setminus X$ *is an essential simple closed curve. If every element of* G *is isotopic rel* X *to a homeomorphism that preserves* γ *and does not interchange the components of* $\mathbb{S}^2 \setminus \gamma$ *then* $\mathrm{Fix}(G) \neq \emptyset$*.*

Proof. Let D be a connected component of $\mathbb{S}^2 \setminus \gamma$. Consider the universal covering Y of the connected component containing D of $\mathbb{S}^2 \setminus (X \setminus D)$. Let $\tilde{D}$ be a lift of D to the topological disc Y. We denote by X_0 the lift of $X \cap D$ to the disc $\tilde{D}$. There exists a unique lift $\tilde{G}$ of G to Y such that $\tilde{\phi}(X_0) = X_0$ for any $\tilde{\phi} \in \tilde{G}$. The set X_0 is compact; there exists a global fixed point for $\tilde{G}$ (and then for G) by Th. 1.3. □

Cartwright and Littlewood [6] proved that an orientation preserving homeomorphism of $\mathbb{R}^2$ that leaves a compact nonseparating continuum (a compact connected set whose complementary is also connected) K invariant leaves a point of K fixed (see [5] for a short proof). We generalize Cartwright-Littlewood theorem for nilpotent subgroups of $\mathrm{Diff}_+^1(\mathbb{S}^2)$.

Proof of Theorem 1.4. Suppose that the result is false. There exists a closed simple curve γ separating K and $\mathrm{Fix}(G) \cup \{\infty\}$ in $\mathbb{S}^2$. Analogously as in Lemma 4.1 we obtain a lift $\tilde{G}$ of G to the universal covering of the connected component of $\mathbb{R}^2 \setminus \mathrm{Fix}(G)$ containing K such that the lift preserves a compact set (a lift of K). Theorem 1.3 implies $\mathrm{Fix}(\tilde{G}) \neq \emptyset$ and then $\mathrm{Fix}(G) \cap (\mathbb{R}^2 \setminus \mathrm{Fix}(G)) \neq \emptyset$. □

Proof of Theorem 1.2. It suffices to show the result for finitely generated subgroups. A finitely generated nilpotent subgroup is polycyclic, so every subgroup is finitely generated (see Theorem 2.7, page 31 [19]). The theorem is proved by induction on the class of nilpotency k of G. It is obvious for $k = 0$. We will prove that if the theorem holds true for groups in $\mathcal{G}_{k,p}$ then then it is also satisfied for groups in $\mathcal{G}_{k,p+1}$. The result is obviously true for $p = 0$ and $p = 1$.

Let $G =< Z^{(k)}(G), \phi_1, \ldots, \phi_{p+1} >$. Our first goal is finding a finite orbit for G. The groups

$$G_1 =< Z^{(k)}(G), \phi_1, \ldots, \phi_p >, \quad G_2 =< Z^{(k)}(G), \phi_{p+1} >$$

are normal in G and have orbits of period at most 2 by the induction hypothesis. We define

$$H_1 =< \phi^2 : \phi \in G_1 >, \quad H_2 =< \phi^2 : \phi \in G_2 > .$$

The group H_j is a characteristic subgroup of G_j and then a normal subgroup of G. Moreover $|G_j : H_j| < \infty$ since G_j is polycyclic (see Lemma 4.1, page 57 [19]) for $j \in \{1, 2\}$. The sets $\mathrm{Fix}(H_1)$ and $\mathrm{Fix}(H_2)$ are non-empty by the existence of orbits of period at most 2 for G_1 and G_2. If $\mathrm{Fix}(H_1) \cap \mathrm{Fix}(H_2) \neq \emptyset$ then $G =< H_1, H_2 >$ has a global fixed point. Hence G has a finite orbit of period at most $|G_1 : H_1||G_2 : H_2|$.

Suppose $\mathrm{Fix}(H_1) \cap \mathrm{Fix}(H_2) = \emptyset$. Let R be the set of reducing curves provided by Proposition 3.1 ($K_1 = \mathrm{Fix}(H_1)$, $K_2 = \mathrm{Fix}(H_2)$). If $R \neq \emptyset$ there exists a finite index normal subgroup H of G fixing all the curves in R and the connected components in $\mathbb{S}^2 \setminus R$ (modulo isotopy). Lemma 4.1 implies that H has a global fixed point, thus G has a finite orbit. If $R = \emptyset$ then $\sharp(\mathrm{Fix}(H_1) \cup \mathrm{Fix}(H_2)) < \infty$ and $\mathrm{Fix}(H_1)$ is a union of finite orbits of G.

Consider a finite G-orbit $\mathcal{O}$. Let us apply Thurston normal form theorem to the group $[G]$ generated by G in the mapping class group of the marked surface $(\mathbb{S}^2, \mathcal{O})$. Consider the tree Γ that is dual to R (see Def. 3.1). The action of G on Γ has a fixed point τ (Prop. 2.1).

If τ is and edge we can consider a normal subgroup H of G of index at most 2 such that H fixes both sides of $\mathbb{S}^2 \setminus \tau$. The group H has a global fixed point by Lemma 4.1. Hence G has a finite orbit of period at most 2.

We can suppose that G fixes a vertex D of Γ. Let $\hat{D}$ be the marked sphere obtained from $\overline{D}$ by contracting the curves in $\overline{D} \cap \mathcal{A}$. Consider $[] : G \to \mathrm{MCG}(G_{|\hat{D}})$; the model for $[G]$ provided by Lemma 2.1 has a normal subgroup L of index at most 2 that fixes 2 points (Remark 2.2).

We claim that the normal subgroup $H := []^{-1}(L)$ of index at most 2 of G has a global fixed point. If L fixes a curve in $\overline{D} \cap \mathcal{A}$ it is a consequence of Lemma 4.1. If L fixes a puncture there is nothing to prove. We can suppose that L fixes an interior point. There exists a lift $\tilde{H}$ of H to the universal covering of $\mathbb{S}^2 \setminus \mathcal{O}$ such that the fixed point set of some element of $\tilde{H}$ is a non-empty compact set by Prop. 2.4. Hence $\tilde{H}$ has a global fixed point by Lemma 3.2 and Theorem 1.3. We obtain a global fixed point for H and a finite orbit of period at most 2 for G. $\square$

5. Fixed-point-free groups

It is clear that the non-existence of a global fixed point for a subgroup G of $\mathrm{Diff}_+^1(\mathbb{S}^2)$ is related to torsion phenomena (see Remark 2.2). So it is interesting to identify properties that hold true for finite nilpotent groups and to extend them for general nilpotent groups. For instance a finite fixed-point-free nilpotent group is a dihedral group D_{2^n} for some $n \in \mathbb{N}$. This group has a 2-orbit, two 2^n-orbits and every other orbit is a 2^{n+1}-orbit. The generalization of this property is provided by Theorem 1.5: every finite orbit of a fixed-point-free nilpotent subgroup of $\mathrm{Diff}_+^1(\mathbb{S}^2)$ has an even number of elements.

Proof of Theorem 1.5. It suffices to prove the theorem for any finitely generated subgroup H of G. The set F is a union of orbits of H, so H has an odd orbit. We can suppose without lack of generality that G is finitely generated and that F is a G-orbit. We also suppose that $\sharp F \geq 3$, otherwise there is nothing to prove.

We apply the Thurston decomposition (Prop. 3.1) to the marked surface $(\mathbb{S}^2, F)$. Consider the dual tree Γ (see Def. 3.1). Either there exists a connected component D of $\mathbb{S}^2 \setminus \mathcal{A}$ that is G-invariant modulo isotopy rel F or there exists an invariant element γ of R such that the components of $\mathbb{S}^2 \setminus R$ are permuted by G. The latter situation is impossible since $\sharp F$ is odd.

Consider the surface $\hat{D}$ obtained by contracting the curves in $\overline{D} \cap \mathcal{A}$. If $D \cap F \neq \emptyset$ then F is a subset of D and there are no reducing curves. In particular we have $\hat{D} = (\mathbb{S}^2, F)$. We denote $\hat{F} = F$. If $D \cap F = \emptyset$ we define $\hat{F}$ as the set of points in $\hat{D}$ corresponding to curves in $\overline{D} \cap \mathcal{A}$. Consider a curve γ in $\overline{D} \cap \mathcal{A}$. The connected component D_γ of $\mathbb{S}^2 \setminus \gamma$ such that $D_\gamma \cap D = \emptyset$ contains $p \geq 2$ points. The action of G on F is transitive, hence p does not depend on γ and $\sharp F = pq$ where $q \geq 3$ is the cardinal of $\hat{F}$. We deduce that $\sharp \hat{F}$ is odd.

Let $[] : G \to \mathrm{MCG}(\hat{D})$ be the canonical projection. Consider a nilpotent model G_0 of $[G]$ whose elements are all irreducible (Lemma 2.1). By construction $\hat{F}$ is an odd orbit of G_0. Let us show that G_0 has a global fixed point. The group G_0 is of the form $< \mathrm{Tor}(G_0), f_0 >$ where $\mathrm{Tor}(G_0)$ is the normal subgroup of finite order elements of G_0 and either $f_0 \equiv Id$ or f_0 is pseudo-Anosov (Lemma 2.1). The existence of a global fixed point is obvious if $\mathrm{Tor}(G_0) = \{Id\}$.

Suppose $\mathrm{Tor}(G_0) \neq \{Id\}$. Thus there exists $h \in \mathrm{Tor}(G_0) \cap Z^{(1)}(G_0)$ with $h \neq Id$. If the two points in $\mathrm{Fix}(h)$ are not fixed points of G_0 then there exists $h' \in G_0$ permuting the points in $\mathrm{Fix}(h)$ (and commuting with h). A theorem of Handel (see Lemma 5.3) implies $h^2 \equiv Id$. The set $\hat{F}$ is a union of orbits of h. Since $\sharp \hat{F}$ is odd then it contains exactly an element of $\mathrm{Fix}(h)$. Clearly it is a global fixed point.

Let us remark that there are no fixed points of G_0 in $\hat{F}$. Since $[G] \neq \{Id\}$ (the action on punctures and reducing curves is not trivial) there exists a lift $\tilde{G}$ of G to the universal covering of $\mathbb{S}^2 \setminus F$ with an element $\tilde{\alpha} \in \tilde{G}$ satisfying that $\mathrm{Fix}(\tilde{\alpha})$ is non-empty and compact (Prop. 2.4). Hence Theorem 1.3 and Lemma 3.2 imply that $\mathrm{Fix}(\tilde{G})$ (and then $\mathrm{Fix}(G)$) is not empty. □

5.1. **Conditions on 2-orbits.** We study the structure of the 2-orbits of a nilpotent subgroup G of $\mathrm{Diff}^1_+(\mathbb{S}^2)$ and no global fixed points.

Let us begin the section with some examples of finite groups. Consider the subgroup $< 1/z, \lambda z >$ of $\mathrm{PGL}(2, \mathbb{C})$ where λ is a primitive 2^nth root of unity. This group is isomorphic to the dihedral group D_{2^n}. Every finite fixed-point-free nilpotent group of orientation preserving homeomorphisms is topologically conjugated to D_{2^n} for some $n \in \mathbb{N}$ (see Remark 2.2). The group D_2 is commutative and has three 2-orbits $\{0, \infty\}$, $\{1, -1\}$, $\{i, -i\}$. The group D_{2^n} is not commutative and has a unique 2-orbit $\{0, \infty\}$ for any $n \geq 2$. We see in this section that nilpotent subgroups of $\mathrm{Diff}_{+}^{1}(\mathbb{S}^2)$ with no global fixed points share analogous properties.

Definition 5.1. Consider 2-orbits $\mathcal{O}_1$ and $\mathcal{O}_2$ of a subgroup G of $\mathrm{Homeo}_{+}(\mathbb{S}^2)$. We say that $[\mathcal{O}_1] = [\mathcal{O}_2]$ if we have

$$\phi_{|\mathcal{O}_1} \equiv Id \Leftrightarrow \phi_{|\mathcal{O}_2} \equiv Id$$

for any $\phi \in G$.

Definition 5.2. Consider 2-orbits $\mathcal{O}_1$, ..., $\mathcal{O}_n$ of a subgroup G of $\mathrm{Homeo}_{+}(\mathbb{S}^2)$. The mapping

$$\begin{array}{rccc} \tau & : & G & \to & (\mathbb{Z}/2\mathbb{Z})^n \\ & & \phi & \mapsto & (a_1, \ldots, a_n) \end{array}$$

where $a_j = 0$ if and only if $\phi_{|\mathcal{O}_j} \equiv Id$ is a morphism of groups. We say that $\mathcal{O}_1$, ..., $\mathcal{O}_n$ are *independent* if the action of G on $\mathcal{O}_1 \cup \ldots \cup \mathcal{O}_n$ is $(\mathbb{Z}/2\mathbb{Z})^n$, i.e. if $\tau(G)$ is equal to $(\mathbb{Z}/2\mathbb{Z})^n$.

Remark 5.1. If the classes of 2-orbits $\mathcal{O}_1$, $\mathcal{O}_2$ are different ($[\mathcal{O}_1] \neq [\mathcal{O}_2]$) then the action of G on $\mathcal{O}_1 \cup \mathcal{O}_2$ is $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} = D_2$.

Remark 5.2. Two different classes of 2-orbits are always independent. This is not the case for 3 classes of 2-orbits. For instance the subgroup $D_2 =< 1/z, -z >$ of $\mathrm{PGL}(2, \mathbb{C})$ has three 2-orbits whose classes are pairwise different but they are not independent since the action of G on the union of the orbits is D_2.

Lemma 5.1. *Consider 2-orbits $\mathcal{O}_1$, $\mathcal{O}_2$, $\mathcal{O}_3$, $\mathcal{O}_4$ of a subgroup G of $\mathrm{Homeo}_{+}(\mathbb{S}^2)$ whose classes are pairwise different. Then there are three of them that are independent.*

Proof. Suppose that there is no subset of 3 independent 2-orbits. We can consider the action of G on $\mathcal{O}_1 \cup \mathcal{O}_2 \cup \mathcal{O}_3 \cup \mathcal{O}_4$ as a subgroup of $(\mathbb{Z}/2\mathbb{Z})^4$. Indeed consider the mapping τ in Definition 5.2 for $n = 4$. If an element a of $\tau(G)$ has 2 coordinates a_j and a_k ($j \neq k$) equal to 0 then any other coordinate a_l of a is equal to 0. Otherwise $\mathcal{O}_j$, $\mathcal{O}_k$ and $\mathcal{O}_l$ are independent. The projection of $\tau(G)$ in any 2 coordinates is equal to $(\mathbb{Z}/2\mathbb{Z})^2$ since different 2-orbits are independent. We deduce that $(0,1,1,1)$ and $(1,0,1,1)$ belong to $\tau(G)$. Hence its sum $(1,1,0,0)$ belongs to $\tau(G)$ providing an element with exactly two vanishing coordinates. ☐

We already know that $D_2 =< 1/z, -z >$ and $D_4 =< 1/z, iz >$ have 3 and 1 classes of 2-orbits respectively. Moreover the unique commutative finite group in $\mathrm{Homeo}_{+}(\mathbb{S}^2)$ with no global fixed point is D_2 that has

3 classes of 2-orbits. These properties are generalized in Theorem 1.6. The next result is a corollary of Theorem 1.6.

Corollary 5.1. *Let* $G \subset \mathrm{Diff}^1_+(\mathbb{S}^2)$ *be a nilpotent group with either exactly* 2 *or at least* 4 *classes of* 2*-orbits. Then* G *has a global fixed point.*

Given ϕ, η commuting elements of $\mathrm{Homeo}_+(\mathbb{S}^2)$ there exist isotopies ϕ_t and η_t such that $\phi_0 = \eta_0 = Id$, $\phi_1 = \phi$, $\eta_1 = \eta$. The invariant $w(\phi, \eta)$ is the homotopy class defined by $\gamma_t = \phi_t^{-1} \eta_t^{-1} \phi_t \eta_t$ in $\pi_1(\mathrm{Homeo}_+(\mathbb{S}^2), Id)$. This group is isomorphic to $\mathbb{Z}/2\mathbb{Z}$ since $\mathrm{Homeo}_+(\mathbb{S}^2)$ is homotopically equivalent to SO(3) (see [14]). Then we can analyze the existence and properties of 2-orbits of commutative groups by using Theorem 1.1. Even if a nilpotent group $G \subset \mathrm{Diff}^1_+(\mathbb{S}^2)$ (with no global fixed point) is not in general commutative we can still apply the properties of the invariant w (see [12] [9]) since its image in a certain mapping class group is abelian.

Lemma 5.2. *Let* $G \subset \mathrm{Diff}^1_+(\mathbb{S}^2)$ *be a fixed-point-free nilpotent group. Let* F *be a finite set composed by two or three* 2*-orbits of* G *whose classes are pairwise different. Then the image* $[G]$ *of* G *by the mapping* $[] : G \to \mathrm{MCG}(\mathbb{S}^2, F)$ *is a commutative irreducible group.*

Proof. We apply the Thurston decomposition (Prop. 3.1) to the marked surface $(\mathbb{S}^2, F)$ with $H_1 = H_2 = G$, $K_1 = \emptyset$, $K_2 = \emptyset$. Suppose that $[G]$ is not irreducible; thus the set R of reducing curves is non-empty. Consider the tree Γ that is dual to R. There is no an invariant element γ of R such that the components of $\mathbb{S}^2 \setminus R$ are permuted by G (modulo isotopy rel F) since such property implies that all the 2-orbits in F are equivalent. Thus there exists a connected component D of $\mathbb{S}^2 \setminus \mathcal{A}$ that is G-invariant modulo isotopy.

We claim that G fixes a curve in $\overline{D} \cap \mathcal{A}$ modulo isotopy. Let D' be a connected component of $\mathbb{S}^2 \setminus \overline{D}$. The set D' contains at least 2 points of F since curves in R are essential. If $F \cap D'$ does not contain a 2-orbit then there are two equivalent 2-orbits in F. This contradicts the choice of F. Hence $F \cap D'$ contains a 2-orbit and $\partial D \cap \partial D'$ is G-invariant modulo isotopy. Lemma 4.1 implies that G has a global fixed point, contradicting the hypothesis.

Consider a nilpotent model G_0 of $[G]$ whose elements are all irreducible (Lemma 2.1). The set $\mathrm{Tor}(G_0)$ of finite order elements is a subgroup of G_0 (Lemma 2.1). It is either a cyclic group C_n or a dihedral group D_{2^n} for some $n \in \mathbb{N}$ (see Remark 2.2). The set F contains at least two disjoint invariant sets of 2 elements of $\mathrm{Tor}(G_0)$. Hence we obtain $\mathrm{Tor}(G_0) = \{Id\}$, $\mathrm{Tor}(G_0) = C_2$ or $\mathrm{Tor}(G_0) = D_2$. In the first case G_0

is clearly commutative. In the second case $Z^{(1)}(G_0) \cap \mathrm{Tor}(G_0) \neq \{Id\}$ implies $\mathrm{Tor}(G_0) \subset Z^{(1)}(G_0)$. Since $G_0/\mathrm{Tor}(G_0)$ is cyclic then G_0 is commutative. We can suppose $\mathrm{Tor}(G_0) = D_2$. The group $\mathrm{Tor}(G_0)$ has exactly three 2-orbits. Each one of them is the fixed point set of one of the three non-trivial elements of $\mathrm{Tor}(G_0)$. Let $h \in \mathrm{Tor}(G_0) \setminus \{Id\}$. If $fhf^{-1} = h' \neq h$ for some $f \in G_0$ then $\mathrm{Fix}(h) \cup \mathrm{Fix}(h')$ is contained in an orbit of G_0 and G_0 has at most an invariant set of 2 elements. We deduce $\mathrm{Tor}(G_0) \subset Z^{(1)}(G_0)$; the group G_0 is commutative. □

Definition 5.3. Consider the setup in Lemma 5.2. We can define an invariant $w_F : G \times G \to \mathbb{Z}/2\mathbb{Z}$. Indeed given $\phi, \eta \in G$ we consider elements $\phi_0, \eta_0 \in \mathrm{Homeo}_+(\mathbb{S}^2)$ such that $[\phi] = [\phi_0]$ and $[\eta] = [\eta_0]$ in $\mathrm{MCG}(\mathbb{S}^2, F)$ and $\phi_0\eta_0 = \eta_0\phi_0$. We define $w_F(\phi, \eta) = w(\phi_0, \eta_0)$.

Remark 5.3. Since the center of the group $\pi_1(\mathbb{S}^2 \setminus F)$ is trivial ($\mathbb{S}^2 \setminus F$ has negative Euler characteristic) then the group $\pi_1(\mathrm{MCG}(\mathbb{S}^2, F), Id)$ is also trivial. In particular the definition of $w_F(\phi, \eta)$ does not depend on the choices of ϕ_0 and η_0 (provided they commute). We can choose ϕ_0 and η_0 in the group G_0 defined in Lemma 5.2.

The next lemmas are used to obtain properties of the invariant w_F. They provide the algebraic structure of the invariant and link the existence of fixed points with its vanishing respectively.

Proposition 5.1. *(Prop. 3.2* [9]*) Let G be a commutative subgroup of* $\mathrm{Homeo}_+(\mathbb{S}^2)$*. The invariant $w : G \times G \to \mathbb{Z}/2\mathbb{Z}$ is symmetric and $w(., g)$ is a morphism of groups for any $g \in G$.*

Lemma 5.3. *(Lemma 1.2* [12]*) Suppose that $\eta : \mathbb{S}^2 \to \mathbb{S}^2$ is a rotation about some axis by $2\pi p/q$ ($p/q \in \mathbb{Q}, 0 < p/q < 1$) and that $\phi \in$* $\mathrm{Homeo}_+(\mathbb{S}^2)$ *commutes with η.*

- *If ϕ interchanges the fixed points z_1 and z_2 of η then $p/q = 1/2$ and $w(\phi, \eta) = 1$.*
- *If $\phi(z_1) = z_1$ and $\phi(z_2) = z_2$ then $w(\phi, \eta) = 0$.*

Proof. The first case is Handel's lemma. The second property is an easy exercise whose proof we include for the sake of clarity.

Let A be the open annulus $\mathbb{S}^2 \setminus \{z_1, z_2\}$. Consider the universal covering $\tilde{A}$ of A and a generator T of the group of covering transformations. There exists an isotopy η_t rel $\{z_1, z_2\}$ such that $\eta_0 = Id$ and $\eta_1 = \eta$. For instance η_t can be a rotation by $2\pi tp/q$. Consider the lift $\tilde{\eta}_t$ of η_t to $\tilde{A}$ such that $\tilde{\eta}_0 = Id$. Since the class of an element in the mapping class group of the twice-punctured sphere is determined by the action on the punctures ($\mathrm{MCG}(S_{0,2}) \approx \mathbb{Z}/2\mathbb{Z}$) we consider an isotopy ϕ_t rel $\{z_1, z_2\}$ such that $\phi_0 = Id$ and $\phi_1 = \phi$. We consider the lift $\tilde{\phi}_t$ of ϕ_t

to $\tilde{A}$ such that $\tilde{\phi}_0 = Id$. We have $\tilde{\eta}^q = T^p$, $\tilde{\phi}T = T\tilde{\phi}$, $\tilde{\eta}T = T\tilde{\eta}$ and $\tilde{\eta}\tilde{\phi} = T^a\tilde{\phi}\tilde{\eta}$ for some $a \in \mathbb{Z}$.

We have

$$T^p\tilde{\phi} = \tilde{\eta}^q\tilde{\phi} = T^{aq}\tilde{\phi}\tilde{\eta}^q = T^{aq}\tilde{\phi}T^p = T^{aq+p}\tilde{\phi} \implies p = aq + p \implies a = 0.$$

The class of the path $\gamma_z : [0,1] \to A$ defined by $\gamma(t) = (\phi_t^{-1}\eta_t^{-1}\phi_t\eta_t)(z)$ is trivial in $\pi_1(A, z)$ for any $z \in A$ since $\tilde{\eta}\tilde{\phi} = \tilde{\phi}\tilde{\eta}$. The path $\phi_t^{-1}\eta_t^{-1}\phi_t\eta_t$ is trivial in $\pi_1(\text{Homeo}_+(\mathbb{S}^2, \{z_1, z_2\}), Id)$ (see Lemma 1.1 of [12]). We deduce $w(\phi, \eta) = 0$. $\square$

Next proposition encapsulates the properties of the invariant w_F that allow to prove Theorem 1.6.

Proposition 5.2. *Let $G \subset \text{Diff}_+^1(\mathbb{S}^2)$ be a fixed-point-free nilpotent group. Let F be a finite set composed by two or three 2-orbits of G whose classes are pairwise different. Then we have*

- *The invariant $w_F : G \times G \to \mathbb{Z}/2\mathbb{Z}$ is symmetric and $w_F(., \eta)$ is a morphism of groups for any $\eta \in G$.*
- *Given a subgroup H of G such that $(w_F)_{|H \times H} \equiv 0$ then H has a global fixed point. In particular we have $w_F \not\equiv 0$.*

Proof. The image $[G]$ of G in $\text{MCG}(\mathbb{S}^2, F)$ is an irreducible commutative group (Lemma 5.2). Let G_0 be the model of $[G]$ provided by Lemma 2.1. Since $w : G_0 \times G_0 \to \mathbb{Z}/2\mathbb{Z}$ is symmetric and a morphism of groups in each component (Proposition 5.1) so is w_F.

Consider the subgroup H_0 of G_0 that is the image of $[H]$ by the isomorphism $[G] \to G_0$ that associates elements defining the same class in $\text{MCG}(\mathbb{S}^2, F)$.

We claim that $\text{Fix}(H_0)$ is not empty. If $\text{Tor}(H_0) = \{Id\}$ then H_0 is cyclic and $\text{Fix}(H_0) \neq \emptyset$. Let $h \in \text{Tor}(H_0) \setminus \{Id\}$. The set $\text{Fix}(h)$ has 2 points and it is H_0-invariant. Lemma 5.3 implies $\text{Fix}(h) = \text{Fix}(H_0)$.

We can suppose $\text{Fix}(H_0) \cap F = \emptyset$ since $\text{Fix}(H_0) \cap F = \text{Fix}(H) \cap F$. There exists a lift $\tilde{H}$ of H to the universal covering of $\mathbb{S}^2 \setminus F$ with an element $\tilde{\alpha} \in \tilde{H}$ such that $\text{Fix}(\tilde{\alpha})$ is a non-empty compact set (Prop. 2.4). Theorem 1.3 and Lemma 3.2 imply that $\text{Fix}(\tilde{H})$ and then $\text{Fix}(H)$ are not empty. $\square$

Consider the hypotheses in Proposition 5.2. In next lemma we see that w_F is completely determined. The description of w_F implies Theorem 1.6 in a straightforward way. Moreover the mapping w_F is simple, for instance w_F factors through a quotient (isomorphic to D_2) of G.

Lemma 5.4. *Let $G \subset \text{Diff}_+^1(\mathbb{S}^2)$ be a fixed-point-free nilpotent group. Let F be a finite set composed by two or three 2-orbits of G whose* *classes are pairwise different. Fix different* 2*-orbits* $\mathcal{O}_1$, $\mathcal{O}_2$ *contained in* F. *We denote*

$$H_j = \{h \in G : h_{|\mathcal{O}_j} \equiv Id\} \text{ for } j \in \{1,2\}.$$

Then $H_1 \cap H_2$ *coincides with* $J := \{\phi \in G : w_F(\phi, \eta) = 0 \ \forall \eta \in G\}$. *Moreover we have*

- *If* F *is composed of two* 2*-orbits then there exists a third class of* 2*-orbits for* G.
- *If* F *is composed of three* 2*-orbits then they are not independent.*

Proof. Consider the image $[G]$ of G in $\mathrm{MCG}(\mathbb{S}^2, F)$. There exists a realization $\sigma : [G] \to G_0$ of $[G]$ as an irreducible group (Lemma 2.1).

The group H_j is a normal subgroup of index 2 of G for $j \in \{1,2\}$. The groups H_1 and H_2 are different since $[\mathcal{O}_1] \neq [\mathcal{O}_2]$. Consider $\eta_1 \in H_2 \setminus H_1$ and $\eta_2 \in H_1 \setminus H_2$. The group $\sigma(H_j)$ has global fixed points in F and $\sigma(H_j)/\mathrm{Tor}(\sigma(H_j))$ is cyclic. Lemma 5.3 implies that $w_{|\sigma(H_j) \times \sigma(H_j)}$ is trivial and so is $(w_F)_{|H_j \times H_j}$.

Consider $\phi \in H_1 \cap H_2$, we have that $w_F(\phi, \alpha_1) = 0$ for any $\alpha_1 \in H_1$ since $\phi \in H_1$ and $w_F(\phi, \eta_1) = 0$ since $\eta_1, \phi \in H_2$. Since $w_F(\phi, .)$ is a morphism of groups and $G =< H_1, \eta_1 >$ we obtain that ϕ belongs to J. The mapping $(w_F)_{|G \times G}$ is not trivial but $(w_F)_{|H_1 \times H_1}$ is. Thus there exists $\beta_1 \in H_1$ such that $w_F(\beta_1, \eta_1) = 1$. Consider the normal subgroup

$$J_1 = \{\eta \in G : w_F(\eta, \beta_1) = 0\}$$

of G. It is clear that $H_1 \subset J_1$, $J_1 \neq G$ and that J_1 has index 2. Thus H_1 and J_1 coincide and J is contained in H_1. Analogously we obtain $J \subset H_2$. We get $J \subset H_1 \cap H_2$ and then $J = H_1 \cap H_2$.

Notice that if $\mathcal{O}_3$ is a third 2-orbit in F the previous proof shows that $J = H_1 \cap H_2 \cap H_3$ and the orbits of F can not be independent.

Suppose that $F = \mathcal{O}_1 \cup \mathcal{O}_2$. The group $G/J = \{Id, [\eta_1], [\eta_2], [\eta_1\eta_2]\}$ is isomorphic to D_2. We define $H_3 =< J, \eta_1\eta_2 >$, it is a normal subgroup of index 2 of G such that $w_F : H_3 \times H_3 \to \mathbb{Z}/2\mathbb{Z}$ is the trivial mapping. Then H_3 has a global fixed point by Proposition 5.2. The group H_3 has no fixed points in F since $\eta_1\eta_2$ does not fix any element of F. Since H_3 is normal of index 2 in G then there exists a 2-orbit $\mathcal{O}_3 \subset \mathrm{Fix}(H_3)$. We have $(\eta_1\eta_2)_{|\mathcal{O}_3} \equiv Id$, $(\eta_1\eta_2)_{|\mathcal{O}_1} \neq Id$ and $(\eta_1\eta_2)_{|\mathcal{O}_2} \neq Id$. Thus we obtain $[\mathcal{O}_1] \neq [\mathcal{O}_3]$ and $[\mathcal{O}_2] \neq [\mathcal{O}_3]$. ☐

Corollary 5.2. *Let* $G \subset \mathrm{Diff}^1_+(\mathbb{S}^2)$ *be a fixed-point-free nilpotent group. Then there are no* 3 *independent* 2*-orbits.*

Proof of Theorem 1.6. The number of classes of 2-orbits is never 2 since the existence of two 2-orbits implies that there is a third one (Lemma

5.4). Moreover such a number can not be greater than 3 by Lemma 5.1 and Corollary 5.2.

Suppose that G is commutative. Let $\mathcal{O}_1$ be a 2-orbit of G. We denote $H_1 = \{\phi \in G : \phi_{|\mathcal{O}_1} \equiv Id\}$. Since $\mathcal{O}_1$ is contained in $\mathrm{Fix}(H_1)$ we deduce that $w : H_1 \times H_1 \to \mathbb{Z}/2\mathbb{Z}$ is the trivial mapping (Theorem 1.1). Theorem 1.1 also implies that $w : G \times G \to \mathbb{Z}/2\mathbb{Z}$ is not trivial. Let $\eta_1 \in G \setminus H_1$ Since H_1 is an index 2 normal subgroup of G and $w(\eta_1, \eta_1) = 0$ there exists $\alpha_1 \in H_1$ such that $w(\alpha_1, \eta_1) = 1$. We define

$$H_2 = \{\phi \in G : w(\phi, \eta_1) = 0\}.$$

The group H_2 is normal in G, moreover it has index 2 since $\alpha_1 \in H_1 \setminus H_2$. Thus $J := H_1 \cap H_2$ is a normal subgroup of G of index 4. Indeed we have $H_2 =< J, \eta_1 >$. Since $w_{|H_1 \times H_1} \equiv 0$ and $w_{|H_2 \times \{\eta_1\}} \equiv 0$ then we deduce $w_{|H_2 \times H_2} \equiv 0$. Hence the set $\mathrm{Fix}(H_2)$ is non-empty (Theorem 1.1). There exists a 2-orbit $\mathcal{O}_2$ of G contained in $\mathrm{Fix}(H_2)$. Since $(\eta_1)_{|\mathcal{O}_2} \equiv Id$ and $(\eta_1)_{|\mathcal{O}_1} \not\equiv Id$ then we obtain $[\mathcal{O}_1] \neq [\mathcal{O}_2]$. The group G has 3 classes of 2-orbits by the first part of the proof. $\square$

Theorems 1.5 and 1.6 for commutative groups admit simple proofs by using only the characterization of commutative subgroups G of $\mathrm{Diff}^1_+(\mathbb{S}^2)$ with global fixed points of [9] (see Theorem 1.1).

Corollary 5.3. *Let* $G \subset \mathrm{Diff}^1_+(\mathbb{S}^2)$ *be a fixed-point-free nilpotent group. Let* J *be the subgroup of elements of* G *fixing every* 2*-orbit pointwise. Then it is a normal subgroup such that* $G/J \approx \mathbb{Z}/2\mathbb{Z}$ *or* $G/J \approx D_2$ *depending on whether there are* 1 *or* 3 *classes of* 2*-orbits.*

REFERENCES

[1] Gilbert Baumslag. *Lecture notes on nilpotent groups.* Regional Conference Series in Mathematics, No. 2. American Mathematical Society, Providence, R.I., 1971.

[2] Joan S. Birman, Alex Lubotzky, and John McCarthy. Abelian and solvable subgroups of the mapping class groups. *Duke Math. J.*, 50(4):1107–1120, 1983.

[3] Christian Bonatti. Un point fixe commun pour des difféomorphismes commutants de S^2. *Ann. of Math. (2)*, 129(1):61–69, 1989.

[4] Christian Bonatti. Difféomorphismes commutants des surfaces et stabilité des fibrations en tores. *Topology*, 29(1):101–126, 1990.

[5] Morton Brown. A short short proof of the Cartwright-Littlewood theorem. *Proc. Amer. Math. Soc.*, 65(2):372, 1977.

[6] M. L. Cartwright and J. E. Littlewood. Some fixed point theorems. *Ann. of Math. (2)*, 54:1–37, 1951. With appendix by H. D. Ursell.

[7] S. Druck, F. Fang, and S. Firmo. Fixed points of discrete nilpotent group actions on S^2. *Ann. Inst. Fourier (Grenoble)*, 52(4):1075–1091, 2002.

[8] John Franks, Michael Handel, and Kamlesh Parwani. Fixed points of abelian actions. *J. Mod. Dyn.*, 1(3):443–464, 2007.

[9] John Franks, Michael Handel, and Kamlesh Parwani. Fixed points of abelian actions on S^2. *Ergodic Theory Dynam. Systems*, 27(5):1557–1581, 2007.

[10] J.-M. Gambaudo. Periodic orbits and fixed points of a C^1 orientation-preserving embedding of D^2. *Math. Proc. Cambridge Philos. Soc.*, 108(2):307–310, 1990.

[11] Étienne Ghys. Sur les groupes engendrés par des difféomorphismes proches de l'identité. *Bol. Soc. Brasil. Mat. (N.S.)*, 24(2):137–178, 1993.

[12] Michael Handel. Commuting homeomorphisms of S^2. *Topology*, 31(2):293–303, 1992.

[13] Steven P. Kerckhoff. The Nielsen realization problem. *Ann. of Math. (2)*, 117(2):235–265, 1983.

[14] Hellmuth Kneser. Die Deformationssätze der einfach zusammenhängenden Flächen. *Math. Z.*, 25(1):362–372, 1926.

[15] Patrice Le Calvez. Pourquoi les points périodiques des homéomorphismes du plan tournent-ils autour de certains points fixes? *Ann. Sci. Éc. Norm. Supér. (4)*, 41(1):141–176, 2008.

[16] Elon L. Lima. Common singularities of commuting vector fields on 2-manifolds. *Comment. Math. Helv.*, 39:97–110, 1964.

[17] Gregory Margulis. Free subgroups of the homeomorphism group of the circle. *C. R. Acad. Sci. Paris Sér. I Math.*, 331(9):669–674, 2000.

[18] J. F. Plante. Fixed points of Lie group actions on surfaces. *Ergodic Theory Dynam. Systems*, 6(1):149–161, 1986.

[19] M. S. Raghunathan. *Discrete subgroups of Lie groups*. Springer-Verlag, New York, 1972. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 68.

[20] Jean-Pierre Serre. *Trees*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. Translated from the French original by John Stillwell, Corrected 2nd printing of the 1980 English translation.

[21] Jerry Shurman. *Geometry of the quintic*. A Wiley-Interscience Publication. John Wiley & Sons Inc., New York, 1997.

INSTITUTO DE MATEMÁTICA, UFF, RUA MÁRIO SANTOS BRAGA S/N VALONGUINHO, NITERÓI, RIO DE JANEIRO, BRASIL 24020-140